

Math 217 Fall 2025

Quiz 20 – Solutions

Dr. Samir Donmazov

1. Complete* the partial sentences below into precise definitions for, or precise mathematical characterizations of, the italicized term:

- (a) Let $\mathfrak{B} = (v_1, \dots, v_d)$ be a basis for the vector space V . Let $v \in V$. The *\mathfrak{B} -coordinates* of v are ...

Solution: The unique scalars $a_1, \dots, a_d \in \mathbb{R}$ such that

$$v = a_1 v_1 + \dots + a_d v_d.$$

- (b) Let $\mathfrak{B} = (v_1, \dots, v_d)$ be a basis for the vector space V . Let $v \in V$. The *\mathfrak{B} -coordinate column vector* of v is ...

Solution: The column vector formed from the \mathfrak{B} -coordinates of v :

$$[v]_{\mathfrak{B}} = \begin{bmatrix} a_1 \\ \vdots \\ a_d \end{bmatrix} \in \mathbb{R}^d, \quad \text{where } v = a_1 v_1 + \dots + a_d v_d.$$

- (c) Let $T : V \rightarrow V$ be linear and $\mathfrak{B} = (v_1, \dots, v_n)$ an ordered basis of V . The *matrix of T with respect to \mathfrak{B}* is ...

Solution: The $n \times n$ matrix $[T]_{\mathfrak{B}}$ whose j -th column is $[T(v_j)]_{\mathfrak{B}}$; i.e.

$$[T]_{\mathfrak{B}} = \begin{bmatrix} | & & | \\ [T(v_1)]_{\mathfrak{B}} & \cdots & [T(v_n)]_{\mathfrak{B}} \\ | & & | \end{bmatrix}.$$

2. Suppose V is a vector space and v_1, \dots, v_m are linearly independent in V . Show that if $v \in V \setminus \text{Span}(v_1, \dots, v_m)$, then the vectors v, v_1, \dots, v_m are linearly independent.

Solution: Suppose $a v + \sum_{i=1}^m b_i v_i = 0$. If $a \neq 0$, then

$$v = -\frac{1}{a} \sum_{i=1}^m b_i v_i \in \text{Span}(v_1, \dots, v_m),$$

*For full credit, please write out fully what you mean instead of using shorthand phrases.

contradiction. Hence $a = 0$, so $\sum b_i v_i = 0$. By linear independence of v_1, \dots, v_m , we get $b_1 = \dots = b_m = 0$. Thus v, v_1, \dots, v_m are linearly independent.

3. True or False. If you answer true, then state TRUE. If you answer false, then state FALSE. Justify your answer with either a short proof or an explicit counterexample.

(a) Suppose $V = \mathbb{R}^{2 \times 2}$ with basis $\mathfrak{B} = \left(\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right)$. Consider $T : \mathbb{R}^{2 \times 2} \rightarrow \mathbb{R}^{2 \times 2}$ given by $T(X) = AX - XA$ where $A = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$. The matrix of T with respect to \mathfrak{B} is

$$\begin{bmatrix} 0 & 0 & 4 & 0 \\ -4 & 0 & 0 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -4 & 0 \end{bmatrix}.$$

Solution: FALSE. Let $E_{11}, E_{12}, E_{21}, E_{22}$ be the listed basis. Compute:

$$T(E_{11}) = AE_{11} - E_{11}A = \begin{bmatrix} 0 & -3 \\ 0 & 0 \end{bmatrix} = -3E_{12},$$

$$T(E_{12}) = AE_{12} - E_{12}A = \mathbf{0},$$

$$T(E_{21}) = AE_{21} - E_{21}A = 3E_{11} - 3E_{22},$$

$$T(E_{22}) = AE_{22} - E_{22}A = 3E_{12}.$$

Thus the columns of $[T]_{\mathfrak{B}}$ are

$$\begin{bmatrix} 0 \\ -3 \\ 0 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 3 \\ 0 \\ 0 \\ -3 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 3 \\ 0 \\ 0 \end{bmatrix},$$

so

$$[T]_{\mathfrak{B}} = \begin{bmatrix} 0 & 0 & 3 & 0 \\ -3 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & -3 & 0 \end{bmatrix},$$

not the given matrix with 4's.